

Total No. of Questions :8]

SEAT No. :

**P4021**

**[5351]-101**

[Total No. of Pages : 4

**F.E.**

**ENGINEERING MATHEMATICS-I**  
**(2015 Pattern) (Semester - I & II) (Credit System)**

*Time : 3 Hours]*

*[Max. Marks : 50*

*Instructions to the candidates:*

- 1) *Solve Q. 1 or Q. 2, Q. 3 or Q. 4, Q. 5 or Q. 6 and Q. 7 or Q. 8.*
- 2) *Neat diagram must be drawn wherever necessary.*
- 3) *Figures to the right indicate full marks.*
- 4) *Use of, electronic pocket calculator allowed.*
- 5) *Assume suitable data, if necessary.*

**Q1) a)** Reduce the following matrix to its normal form and hence find its rank.

**[4]**

$$A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix}$$

b) Find the eigen values and eigen vectors of :

**[4]**

$$A = \begin{bmatrix} -1 & 1 & 2 \\ 0 & -2 & 1 \\ 0 & 0 & -3 \end{bmatrix}$$

c) If  $\tan(\alpha + i\beta) = x + iy$ , then prove that  $x^2 + y^2 + 2x \cot 2\alpha = 1$  and  $x^2 + y^2 - 2y \coth(2\beta) + 1 = 0$ .

**[4]**

OR

**Q2) a)** If  $\alpha$  and  $\beta$  are roots of equation  $Z^2 \sin^2 \theta - Z \sin 2\theta + 1 = 0$ , prove that

$$\alpha^n + \beta^n = 2 \cos n\theta \cdot \operatorname{cosec}^n \theta, \text{ where } n \text{ is integer.}$$

**[4]**

b) Prove that  $\tan \left\{ i \log \left( \frac{a-ib}{a+ib} \right) \right\} = \frac{2ab}{a^2 - b^2}$  using principal value of logarithm.

**[4]**

c) Examine Linear dependence of vectors  $x_1 = (2, -1, 3, 2)$ ,  $x_2 = (1, 3, 4, 2)$ ,  $x_3 = (-1, -4, 1, 0)$ . If dependent find the relation among them.

**[4]**

**P.T.O.**

**Q3) a)** Solve any one : **[4]**

i) Test the convergence of the series  $\sum_{n=1}^{\infty} \frac{(n+1)x^n}{n^2}$ .

ii) Test the convergence of the series  $1 - \frac{1}{3} + \frac{1}{9} - \frac{1}{27} + \dots$

b) Expand  $3x^3 - 2x^2 + x - 4$  in powers of  $(x+2)$  using Taylor's theorem. **[4]**

c) Find the  $n^{\text{th}}$  derivative of  $y = \cos^{-1}\left(\frac{x - x^{-1}}{x + x^{-1}}\right)$ . **[4]**

**Q4) a)** Solve any one : **[4]**

i)  $\lim_{x \rightarrow 0} \left( \frac{2^x + 3^x}{2} \right)^{1/x}$

ii)  $\lim_{x \rightarrow 0} \frac{xe^x - \log(1+x)}{x^2}$

b) Show that : **[4]**

$$(1+x)^x = 1 + x^2 - \frac{1}{2}x^3 + \frac{5}{6}x^4 - \frac{3}{4}x^5 + \dots$$

c) If  $y = e^{\tan^{-1}x}$ , then prove that

$$(1+x^2)y_{n+2} + [2(n+1)x - 1]y_{n+1} + n(n+1)y_n = 0$$
 **[4]**

**Q5) Solve any two :**

a) If  $u = 2x + 3y$ ,  $v = 3x - 2y$  find the value of  $\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v \left(\frac{\partial y}{\partial v}\right)_x \left(\frac{\partial v}{\partial y}\right)_u$ . **[6]**

b) If  $u = \operatorname{cosec}^{-1} \left( \sqrt{\frac{x^{1/2} + y^{1/2}}{x^{1/3} + y^{1/3}}} \right)$ , show that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \frac{\tan u}{12} \left( \frac{13}{12} + \frac{\tan^2 u}{12} \right). \quad [7]$$

c) If  $z = f(x, y)$  where  $x = u + v$ ,  $y = uv$ , then prove that

$$u \frac{\partial z}{\partial u} + v \frac{\partial z}{\partial v} = x \frac{\partial z}{\partial x} + 2y \frac{\partial z}{\partial y} \quad [6]$$

OR

**Q6)** Solve any two :

a) If  $u = \log(x^3 + y^3 - x^2y - xy^2)$ , prove that  $\left( \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)^2 u = \frac{-4}{(x+y)^2}$ . [6]

b) If  $u = \frac{x^3 + y^3}{x + y} + \frac{1}{x^5} \sin^{-1} \left[ \frac{x^2 + y^2}{x^2 + 2xy} \right]$ , find the value of

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} \text{ at the point } (1, 2). \quad [7]$$

c) If  $u = x^2 - y^2$ ,  $v = 2xy$  and  $z = f(u, v)$  then show that

$$x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y} = 2\sqrt{u^2 + v^2} \frac{\partial z}{\partial u}. \quad [6]$$

**Q7) a)** If  $ux + vy = a$ ,  $\frac{u}{x} + \frac{v}{y} = 1$ , prove that  $\left( \frac{\partial u}{\partial x} \right)_y - \left( \frac{\partial v}{\partial y} \right)_x = \frac{x^2 + y^2}{y^2 - x^2}$ . [4]

b) In calculating the volume of a right circular cone, errors of 2% and 1% are made in measuring the height and radius base respectively. Find the error in the calculated volume. [4]

- c) Find the stationary points of the function

$f(x, y) = x^3 + 3xy^2 - 15x^2 - 15y^2 + 72x$ . Examine for maxima and minima at these points. [5]

OR

- Q8) a) If  $u = x + y^2, v = y + z^2, w = z + x^2$  find  $\frac{\partial x}{\partial u}$ . [4]

- b) If  $u = x + y + z, v = x^2 + y^2 + z^2, w = xy + yz + zx$ . Examine whether  $u, v, w$  are functionally dependent. If so find the relation between them. [4]

- c) Find the stationary value of  $u = x^m y^n z^p$  under the condition  $x + y + z = a$ . [5]

